

**WEST**[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)**Search Results -**

Terms	Documents
L7 and (residual activity)	21

Database: 

US Patents Full-Text Database	▲
JPO Abstracts Database	
EPO Abstracts Database	
Derwent World Patents Index	
IBM Technical Disclosure Bulletins	▼

Refine Search: 

L7 and (residual activity)	▲
	▼

[Clear](#)

**Search History****Today's Date: 10/16/2000**

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT,JPAB,EPAB,DWPI	L7 and (residual activity)	21	<u>L9</u>
USPT,JPAB,EPAB,DWPI	L7 and (pH adj 6.0)	38	<u>L8</u>
USPT,JPAB,EPAB,DWPI	L1 and (animal adj feed)	167	<u>L7</u>
USPT,JPAB,EPAB,DWPI	L5 and (residual adj activity)	16	<u>L6</u>
USPT,JPAB,EPAB,DWPI	L4 and (pH adj 6.0)	61	<u>L5</u>
USPT,JPAB,EPAB,DWPI	L1 and neutral	258	<u>L4</u>
USPT,JPAB,EPAB,DWPI	L2 and (pH adj 6.0)	22	<u>L3</u>
USPT,JPAB,EPAB,DWPI	L1 and (residual adj activity)	57	<u>L2</u>
USPT,JPAB,EPAB,DWPI	xylanase	1213	<u>L1</u>

**NCBI PubMed PubMed QUERY PubMed ?**Other Formats: **Citation** **MEDLINE**Links: **Related Articles**☐ Order this document*Appl Microbiol Biotechnol* 1993 Dec;40(4):508-14

## **A comparison of two xylanases from the thermophilic fungi *Thielavia terrestris* and *Thermoascus crustaceus*.**

**Gilbert M, Yaguchi M, Watson DC, Wong KK, Breuil C, Saddler JN**

University of Ottawa, Department of Biology, Ontario, Canada.

Two thermophilic xylanases (xylanase II from *Thielavia terrestris* 255B and the 32-kDa xylanase from *Thermoascus crustaceus* 235E) were studied to determine if they had different and complementary modes of action when they hydrolysed various types of xylans. Partial amino acid sequencing showed that these two enzymes belonged to different families of beta-1,4-glycanases. Xylanase II achieved faster solubilization of insoluble xylan whereas the 32-kDa xylanase was more effective in producing xylose and short xylo-oligomers. An assessment of the combined hydrolytic action of the two xylanases did not reveal any co-operative action. The sugars released when the two thermophilic xylanases were used together were almost identical to those released when the 32-kDa xylanase acted alone. The two xylanases were able to remove about 12% of the xylan remaining in an aspen kraft pulp. This indicated that either one of these thermophilic enzymes may be useful for enhancing the bleaching of kraft pulps.

PMID: 7764332, UI: 94114174

 the above report in   format documents on this page through Loansome Doc